



TITLE:

Inner segment ellipsoid band length is a prognostic factor in retinitis pigmentosa associated with EYS mutations: 5-year observation of retinal structure

AUTHOR(S):

Miyata, M; Ogino, K; Gotoh, N; Morooka, S;
Hasegawa, T; Hata, M; Yoshimura, N

CITATION:

Miyata, M ...[et al]. Inner segment ellipsoid band length is a prognostic factor in retinitis pigmentosa associated with EYS mutations: 5-year observation of retinal structure. Eye 2016, 30: 1588-1592

ISSUE DATE:

2016-08-26

URL:

<http://hdl.handle.net/2433/226161>

RIGHT:

This is the accepted version of the article, which has been published in final form at <http://doi.org/10.1038/eye.2016.196>. The full-text file will be made open to the public on 26 February 2017 in accordance with publisher's 'Terms and Conditions for Self-Archiving'. ; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。 ; This is not the published version. Please cite only the published version.

1 What was known before:

2 The inner segment ellipsoid band is correlated with visual function. EYS is an important and common cause of
3 retinitis pigmentosa.

4

5

6 What this study adds:

7 The inner segment ellipsoid band of retinitis pigmentosa patients with EYS mutations shortened during the 5
8 years of observation annually. The length of the inner segment ellipsoid band is a sensitive prognostic factor
9 for the rate of ISe shortening in RP patients with EYS mutations.

10

11 Inner Segment Ellipsoid Band Length is a Prognostic factor in Retinitis
12 Pigmentosa Associated with *EYS* mutations: 5-year Observation of Retinal
13 Structure

14
15 Manabu Miyata, M.D., Ph.D., Ken Ogino, M.D., Ph.D, Norimoto Gotoh, M.D., Ph.D, Satoshi
16 Morooka, M.D., Ph.D., Tomoko Hasegawa, M.D., Masayuki Hata, M.D., Nagahisa
17 Yoshimura, M.D., Ph.D.

18
19 Department of Ophthalmology and Visual Sciences, Kyoto University Graduate School of
20 Medicine, Kyoto, Japan

21
22 Corresponding author:

23 Manabu Miyata, M.D., Ph.D.

24 Department of Ophthalmology and Visual Sciences, Kyoto University Graduate School of
25 Medicine, Shogoin Kawahara-cho 54, Sakyo-ku, Kyoto, 606-8507, Japan

26 TEL: +81-75-751-3248

27 FAX: +81-75-752-0933

28 Email: miyatam@kuhp.kyoto-u.ac.jp

29
30 Running title:

31 Prognostic factor of retinitis pigmentosa

32
33 Key words:

34 EYS; inner segment ellipsoid band; optical coherence tomography; retinitis pigmentosa

35
36 Conflict of Interest statement:

37 The authors declare no competing financial interests.

38
39 Funding/Support:

40 This work was partly supported by the Innovative Techno-Hub for Integrated Medical
41 Bio-Imaging of the Project for Developing Innovation Systems, from the Ministry of
42 Education, Culture, Sports, Science and Technology (MEXT), Japan.

43
44 Financial Disclosures:

45 N. Yoshimura: Topcon Corporation, Tokyo, Japan (Financial Support), Nidek, Gamagori,
46 Japan (Financial Support, Consultant), Canon, Tokyo, Japan (Financial Support).

Abstract**PURPOSE:**

To evaluate whether the length of the inner segment ellipsoid band (ISe) can be used as a prognostic factor for disease course in retinitis pigmentosa (RP) patients with *EYS* mutations by observation over a period of 5 years.

METHODS:

Twelve RP patients with *EYS* mutations were studied. The horizontal and vertical ISe length of the right eye was manually measured at five time-points annually, using spectral domain optical coherence tomography. A regression line through the five points from baseline to the final measurement was drawn and the ratio of the length (%) at each point to the baseline length was calculated; the slope was defined as the rate of ISe shortening (%/year). The correlation between rate of ISe shortening and age, visual acuity, and mean deviation (MD) value were evaluated. The intraclass correlation coefficient (ICC) for the measurements was calculated.

RESULTS:

The mean rate of ISe shortening was $-4.65 \pm 2.89\%$ per year, and the decline was statistically significant. The rate of shortening was significantly negatively correlated with the baseline length ($P = 0.046$, $r = 0.58$), but not with the baseline age, visual acuity, and MD value. The ICC (2, 1) was 0.999.

CONCLUSION:

ISe of all RP patients with *EYS* mutations shortened during the 5 years of annual observation. The measurement of the length of ISe is simple and convenient method with high repeatability and the length is a sensitive prognostic factor for the rate of ISe shortening in RP patients with *EYS* mutations.

Introduction

Retinitis pigmentosa (RP), a set of retinal diseases featuring degeneration of rod and cone photoreceptors, varies with regard to the onset of symptoms, inheritance mode, fundus appearance, and prognosis,¹ possibly because of differences in causative gene mutations.^{2,}

³ Sixty RP causative gene mutations were found.⁴

EYS is an important and common cause of RP in the Japanese, Spanish, British, Chinese, Israelis, and Palestinians.⁵⁻⁹ Furthermore, a report has described that *EYS*-associated RP patients share a relatively uniform phenotype with near-normal central visual function up to their 20s.¹⁰ We have previously reported that severity of RP patients with *EYS* mutations was relatively moderate among RP patients with various mutations.¹¹ Thus, we encounter RP patients with *EYS* mutations at relatively high frequency in daily clinical consultation except detection for causative gene mutations, and RP patients with *EYS* mutations have representative feature in RP. It is significant to investigate RP patients with *EYS* mutations.

Although a change in the retinal structure of RP patients with the same genetic mutations over the medium-term (2 years) has been reported,^{12, 13} no study, to our knowledge, has investigated the changes over the longer term (5 years). Because the change of retinal structure is gradual, it is necessary to assess the change using a long-term follow-up data. Moreover, it is important to understand the change when explaining the disease course to patients in clinical practice.

It has been reported that evaluation of changes in the inner segment ellipsoid band (ISe) is useful for the assessment of retinal health, which is correlated with visual function.¹⁴⁻¹⁷ A previous 2-year study found that ISe decreases year-by-year in RP patients.¹⁸

From these perspectives, we conducted this study to evaluate whether the change of length of ISe over a long-term period in RP patients with *EYS* mutations can be used as a prognostic factor in predicting the disease course.

Subjects and Methods

This study was approved by the ethics committee of Kyoto University Graduate School of Medicine (Kyoto, Japan). All study protocols adhered to the tenets of the Declaration of Helsinki. The nature of the study and the possible risks and benefits of participation were explained to all study candidates. All subjects choosing to participate provided written informed consent.

Subjects

We performed gene analyses for 329 Japanese RP patients who visited the Department of Ophthalmology and Visual Sciences, Kyoto University Graduate School of Medicine, Kyoto,

Japan between January 2011 and December 2012 and agreed to provide peripheral blood samples.⁵ All patients underwent comprehensive ophthalmological examinations, including measurement of the best-corrected visual acuity (BCVA) using a decimal visual acuity chart (Landolt chart), indirect ophthalmoscopy, slit-lamp biomicroscopy, SD-OCT (Spectralis HRA+OCT, Heidelberg Engineering, Heidelberg, Germany); mean deviation (MD) value at baseline calculated using a Humphrey field analyzer (HFA; 10-2 SITA Standard Program; Carl Zeiss Meditec, Jena, Germany), and 30-Hz flicker electroretinography (ERG) were also performed. ERG results were recorded according to the International Society for Clinical Electrophysiology of Vision standard protocol recommended in 2008 using LS-C (Mayo Co., Nagoya, Japan) and Neuropack MEB-2204 systems (Nihon Kohden, Tokyo, Japan). All BCVA data were converted to the logarithm of the minimal angle of resolution (logMAR) for statistical analyses. Retinal specialists diagnosed RP using comprehensive ophthalmological examinations. In all the patients, *EYS* mutations were detected by next-generation sequencing. Inclusion criteria were: available SD-OCT images obtained over a period of 5 years, at five time points or more, on different days. Patients for whom ISe could not be detected on OCT images including dotted lines and those who had undergone intraocular surgery during the study period were excluded. Two investigators (MM and TH) determined whether the images were assessable. Those who had undergone intraocular surgery during the study period were also excluded.

Inner Segment Ellipsoid Band Analysis

The length of ISe on horizontal and vertical OCT B-scan crossing the fovea of the right eye was measured over 5 years using SD-OCT images of 30 × 30° scans (approximately 9 × 9 mm), at five time points. The length of a healthy eye is approximately 9000 μm, because ISe is not cut within range of the SD-OCT images. Representative measurements were used for analysis, by averaging horizontal and vertical measurements. Measurements were performed obtained in a random order using the built-in measurement scale provided in the SD-OCT software (Figure 1). Information pertaining to the date of measurement was masked. If the OCT images were obtained at more than five time points, we selected five images obtained at almost equal time intervals. We plotted this data, with the baseline date up to the final measurement date on the x-axis and the percentage the length of ISe relative to that at baseline (%) on the y-axis. A regression line through the five data points from baseline to the final measurement date was then drawn; the slope was defined as the rate of ISe shortening (%/day). To improve comprehensibility, we converted the rate of ISe shortening (%/day) to the rate of ISe shortening (%/year) by multiplying by 365. To assess the repeatability of measurements, another investigator measured the length of ISe of five patients selected at random in the same manner.

Statistical Analysis

Data are presented as the mean \pm standard deviation where applicable. All statistical analyses were performed using SPSS version 21 (IBM, New York, NJ, USA). Student's *t*-tests were used to compare different data sets. Correlations were analyzed using Pearson's correlation coefficients. Linear regression analysis was performed to calculate the rate of ISe shortening. The intraclass correlation coefficient (ICC) value for the length of ISe measurements recorded by the two investigators (MM and TH) was calculated to determine the reliability of measurements. A *P*-value of < 0.05 was considered statistically significant.

Results

In total, 19 patients met the inclusion criteria for the study. Among these, ISe could not be detected in seven. Eventually, 12 patients were included in our analysis. In one patient, MD value at baseline was not available. Table 1 shows the patient characteristics. The observation period was 5.93 ± 0.74 years. The ICC (2, 1) value for the length of ISe measurements was 0.999.

Figure 2 shows the relationship between the length of ISe at each time point, relative to that at baseline, and the follow-up duration. The rate of ISe shortening was $-4.65 \pm 2.89\%$ per year. There was a significant difference between the derived rate of ISe shortening value and 0 ($P < 0.001$), indicating a significant decrease in the length of ISe. When the cases were analyzed individually, there were significant differences between the rate of ISe shortening and 0 in all patients. When we separated the horizontal and vertical measurements to analyze, the rate of horizontal and vertical ISe shortening was -5.17 ± 3.10 and $-4.26 \pm 2.98\%$ per year, respectively. There was no differences in the rate between them ($P = 0.70$).

The rate of ISe shortening and the length of ISe at baseline were significantly correlated ($P = 0.046$, $r = 0.58$, Figure 3), i.e., the rate of ISe shortening value was high in patients with a small length of ISe value at baseline. However, there was no significant correlation between the rate of ISe shortening and age, log MAR visual acuity, and MD value at baseline ($P = 0.84$, 0.30 , and 0.25 , respectively, Figure 3).

The decrease in the MD value varied; the mean decrease was 3.09 ± 3.62 dB in the same 5-year period. Among 12 patients, six decreased by more than 2 dB and six decreased by less than 2 dB. The rate of ISe shortening was -5.17 ± 4.97 and $-3.52 \pm 1.17\%$ per year, respectively. There were no significant differences of the rate between in the two groups ($P = 0.56$).

Discussion

This study evaluated whether the length of ISe, which is correlated with visual function, changes over a long-term period in RP patients with *EYS* mutations for the first

time. The length of ISe of the most RP patients with *EYS* mutations shortened during the 5 years of annual observations and the rate of the length of ISe shortening was significantly higher in patients with a short length of ISe at baseline.

The length of ISe is a sensitive prognostic factor for the rate of ISe shortening in RP patients with *EYS* mutations. However, age, log MAR visual acuity, and MD value were not statistically significant prognostic factors in the present study, although a previous study has reported that the extent of visual fields constriction seemed to correlate better with age than with visual acuity.¹⁰ These findings help to explain the disease course to patients in clinical practice; i.e., disease progression of RP patients with *EYS* mutations with relatively longer ISe should be slow, while that of the patients with relatively shorter ISe should be rapid. However, we cannot predict the degree of disease progression from age, visual acuity, and visual field. To expand the application of this approach, further studies in degenerative retinal diseases with other mutations are needed.

In this study, the rate of ISe shortening varied among patients with the same type of gene mutation, indicating that RP with the same gene mutation may not follow the same course in all patients. This finding is consistent with that in a previous study where patients with *PRPH2* mutations exhibited different phenotypes.¹⁹ The previous studies showed a wide range of phenotypic expression from the same mutation: central areolar choroidal dystrophy, autosomal dominant RP, adult vitelliform macular dystrophy, and cone-rod dystrophy.

Curcio et al showed that cone density decreased steeply with increasing eccentricity, while rod density increased with increasing eccentricity, and cone and rod density became the same at 0.5 mm from the fovea according to their published figure.²⁰ Thus, an ISe shorter than 1000 μm was mainly constituted of cone cells. The results of the present study suggested that cone cells were more easily disordered than rod cells. There is a need for longitudinal study of the decrease in photoreceptors in RP patients using an adaptive optics scanning laser ophthalmoscope.

In the present study, we selected the right eye in all cases to prevent selection bias. However, the symmetry of progression of RP is also of interest. The ISe of left eye at baseline was $3180 \pm 2348 \mu\text{m}$, which was not significantly different from that of the right eye ($3137 \pm 2348 \mu\text{m}$, $P = 0.67$). The ISe of the left eye at final measurement was $2558 \pm 2325 \mu\text{m}$, which was also not significantly different from that of the right eye ($2456 \pm 2292 \mu\text{m}$, $P = 0.20$). Thus, progression of RP in patients with *EYS* mutations was symmetric in the present study.

It is necessary to follow RP patients for at least 4 years to determine the true extent of changes, because RP is a disease with a long course. Indeed, because RP shows a definite but slow decline in visual function and degeneration of retinal structure, large clinical trials have set study periods for longer than 4 years, although their outcomes were different

224 from those of this study.²¹⁻²³

225 We converted the length of ISe to rate (%) in order to exclude the effect of the length
226 of ISe, because this study included patients with various stages of the disease. For instance,
227 although a change in the length of ISe from 8000 μ m to 7000 μ m indicates the same extent of
228 progression as a reduction from 1500 μ m to 500 μ m, the implications are quite different. In
229 fact, ISe at baseline was ranged from 481 to 7589 μ m.

230 This study had some limitations. First, measurement of the length of ISe was manual.
231 Because ISe may not have a clear-cut edge, the measurement of ISe length can incur errors.
232 We attempted to minimize the level of error as much as possible by two means. One was to
233 make use of the slope of the regression line derived from OCT images obtained at five time
234 points over 5 years; analysis of only two data points, i.e., baseline and approximately 5 years,
235 would generate misleading results. The other was that the measurements of the length of
236 ISe showed high repeatability; the ICC (2, 1) value was 0.999, indicating that almost the
237 same length of ISe could be measured by any examiner. The measurement of the length of
238 ISe is thus a relatively simple and facile, but accurate, method. Second, there were few
239 young patients in this study. Although there was no significant correlation between the rate of
240 ISe shortening and age at baseline, the rate of ISe shortening could not be determined in
241 seven of the 19 patients (37%) in the *EYS*-RP group because of absence of assessable ISe,
242 whereas it could be detected when they were younger. If the results of these patients were
243 included in their youth, the results of this study may be different. Suto et al reported that ISe
244 was absent in 40% RP patients with *EYS* mutations.¹⁰ The results of the present study (37%)
245 were consistent with those of that previous report. Third, few patients had an intermediate
246 length ISe at baseline. Among 12 RP patients, nine had ISe less than 4000 μ m, two had ISe
247 more than 6000 μ m, and one had ISe ranged 4000–6000 μ m. Further research of RP
248 patients with middle ISe should be performed.

249 In conclusion, ISe of all RP patients with *EYS* mutations shortened during the 5
250 years of observation annually. The measurement of the length of ISe is simple and
251 convenient method with high repeatability and this length is a sensitive prognostic factor for
252 the rate of ISe shortening in RP patients with *EYS* mutations.

REFERENCES

1. Hartong DT, Berson EL, Dryja TP. Retinitis pigmentosa. *Lancet*. 2006;**368**:1795-1809.
2. Kelsell RE, Gregory-Evans K, Payne AM, Perrault I, Kaplan J, Yang RB, et al. Mutations in the retinal guanylate cyclase (RETGC-1) gene in dominant cone-rod dystrophy. *Hum Mol Genet*. 1998;**7**:1179-1184.
3. Hayward C, Shu X, Cideciyan AV, Lennon A, Barran P, Zareparsa S, et al. Mutation in a short-chain collagen gene, CTRP5, results in extracellular deposit formation in late-onset retinal degeneration: a genetic model for age-related macular degeneration. *Hum Mol Genet*. 2003;**12**:2657-2667.
4. Xu Y, Guan L, Shen T, Zhang J, Xiao X, Jiang H, et al. Mutations of 60 known causative genes in 157 families with retinitis pigmentosa based on exome sequencing. *Hum Genet*. 2014;**133**:1255-1271.
5. Oishi M, Oishi A, Gotoh N, Ogino K, Higasa K, Iida K, et al. Comprehensive molecular diagnosis of a large cohort of Japanese retinitis pigmentosa and Usher syndrome patients by next-generation sequencing. *Invest Ophthalmol Vis Sci*. 2014;**55**:7369-7375.
6. Arai Y, Maeda A, Hiramami Y, Ishigami C, Kosugi S, Mandai M, et al. Retinitis pigmentosa with EYS mutations is the most prevalent inherited retinal dystrophy in Japanese populations. *J Ophthalmol* **2015**;819760.
7. Barragán I, Borrego S, Pieras JI, González-del Pozo M, Santoyo J, et al. Mutation spectrum of EYS in Spanish patients with autosomal recessive retinitis pigmentosa. *Hum Mutat* 2010;**31**:E1772-E1800.
8. Abd El-Aziz MM, O'Driscoll CA, Kaye RS, Barragan I, El-Ashry MF, Borrego S, et al. Identification of novel mutations in the ortholog of Drosophila eyes shut gene (EYS) causing autosomal recessive retinitis pigmentosa. *Invest Ophthalmol Vis Sci* 2010;**51**:4266-4272.
9. Bandah-Rozenfeld D, Littink KW, Ben-Yosef T, Strom TM, Chowers I, Collin RW, et al. Novel null mutations in the EYS gene are a frequent cause of autosomal recessive retinitis pigmentosa in the Israeli population. *Invest Ophthalmol Vis Sci* 2010;**51**:4387-4394.
10. Suto K, Hosono K, Takahashi M, Hiramami Y, Arai Y, Nagase Y, et al. Clinical phenotype in ten unrelated Japanese patients with mutations in the EYS gene. *Ophthalmic Genet*. 2014;**35**:25-34.
11. Ogino K, Oishi A, Oishi M, Gotoh N, Morooka S, Sugahara M, et al. Efficacy of Column Scatter Plots for Presenting Retinitis Pigmentosa Phenotypes in a Japanese Cohort. *Transl Vis Sci Technol*. 2016;**4**:4.
12. Birch DG, Locke KG, Wen Y, Locke KI, Hoffman DR, Hood DC. Spectral-domain optical coherence tomography measures of outer segment layer progression in patients with X-linked retinitis pigmentosa. *JAMA Ophthalmol*. 2013;**131**:1143-1150.
13. Birch DG, Locke KG, Feliuss J, Klein M, Wheaton DK, Hoffman DR, et al. Rates of decline in regions of the visual field defined by frequency-domain optical coherence tomography in

- 291 patients with RPGR-mediated X-linked retinitis pigmentosa. *Ophthalmology*.
- 292 2015;**122**:833-839.
- 293 14. Sandberg MA, Brodkhurst RJ, Gaudio AR, Berson EL. The association between visual
- 294 acuity and central retinal thickness in retinitis pigmentosa. *Invest Ophthalmol Vis*
- 295 *Sci*. 2005;**46**:3349-3354.
- 296 15. Murakami T, Akimoto M, Ooto S, Suzuki T, Ikeda H, Kawagoe N, et al. Association
- 297 between abnormal autofluorescence and photoreceptor disorganization in retinitis
- 298 pigmentosa. *Am J Ophthalmol*. 2008;**145**:687-694.
- 299 16. Aizawa S, Miyamura Y, Baba T, Hagiwara A, Ogata K, Yamamoto S. Correlation
- 300 between visual function and photoreceptor inner/outer segment junction in patients with
- 301 retinitis pigmentosa. *Eye*. 2009;**23**:304-308.
- 302 17. Rangaswamy NV, Patel HM, Locke KG, Hood DC, Birch DG. A comparison of visual field
- 303 sensitivity to photoreceptor thickness in retinitis pigmentosa. *Invest Ophthalmol Vis Sci*.
- 304 2010;**51**:4213-4219.
- 305 18. Hood DC, Ramachandran R, Holopigian K, Lazow M, Birch DG, Greenstein VC. Method
- 306 for deriving visual field boundaries from OCT scans of patients with retinitis pigmentosa.
- 307 *Biomed Opt Express*. 2011;**2**:1106-1114.
- 308 19. Renner AB, Fiebig BS, Weber BH, Wissinger B, Andreasson S, Gal A, et al. Phenotypic
- 309 variability and long-term follow-up of patients with known and novel PRPH2/RDS gene
- 310 mutations. *Am J Ophthalmol*. 2009;**147**:518-530.
- 311 20. Curcio CA, Sloan KR, Kalina RE, Hendrickson AE. Human photoreceptor topography. *J*
- 312 *Comp Neurol*. 1990;**292**:497-523.
- 313 21. Berson EL, Rosner B, Sandberg MA, Hayes KC, Nicholson BW, Weigel-DiFranco C, et
- 314 al. A randomized trial of vitamin A and vitamin E supplementation for retinitis pigmentosa.
- 315 *Arch Ophthalmol*. 1993;**111**:761-772.
- 316 22. Berson EL, Rosner B, Sandberg MA, Weigel-DiFranco C, Moser A, Brockhurst RJ, et al.
- 317 Clinical trial of docosahexaenoic acid in patients with retinitis pigmentosa receiving vitamin A
- 318 treatment. *Arch Ophthalmol*. 2004;**122**:1297-1305.
- 319 23. Berson EL, Rosner B, Sandberg MA, Weigel-DiFranco C, Brockhurst RJ, Hayes KC, et
- 320 al. Clinical trial of lutein in patients with retinitis pigmentosa receiving vitamin A. *Arch*
- 321 *Ophthalmol*. 2010;**128**:403-411.

322 TITLES AND LEGENDS TO FIGURES

323


324 Figure 1. Spectral domain optical coherence tomography images

325 The length of the inner segment ellipsoid band crossing the fovea was measured in a
326 random order using the measurement scale provided in the spectral-domain optical
327 coherence software. Information pertaining to the date of measurement and type of gene
328 was masked.

329

330 Figure 2. Relationship between the percentage length of the inner segment ellipsoid band at
331 the observation point relative to that at baseline and the follow-up duration.

332 The mean \pm standard deviation rate of the inner segment ellipsoid band shortening is -4.65
333 $\pm 2.89\%$ per year.






Table 1. Baseline characteristics of retinitis pigmentosa patients with *EYS* mutations included in this study

Characteristics	Values
Number of patients	12
Patient age (years)	
Mean \pm SD	45.6 \pm 8.6
Range	34 to 63
Patient sex, no. (%)	
Male	3 (25)
Female	9 (75)
Visual acuity in the right eye, logMAR	
Mean \pm SD	0.040 \pm 0.143
Range	-0.176 to 0.222
Axial length in the right eye (mm)	
Mean \pm SD	24.51 \pm 0.99
Range	22.7 to 25.9
Mean deviation value as per Humphrey 10-2 visual field analysis (dB) (n=11)	
Mean \pm SD	-15.12 \pm 7.16
Range	-25.36 to -1.11
The length of the inner segment ellipsoid band measurement (μ m)	
Mean \pm SD	3137 \pm 2350
Range	481 to 7589
Pseudophakia, no. (%)	1 (8)

SD: standard deviation, logMAR: logarithm of minimal angle resolution